



#0900

Wintafect

DISINFECTANT, CLEANER, & DEODORIZER

DESCRIPTION:

WINTAFECT is a proven disinfectant, cleaner, fungicide, mildewstat, and virucide* with a neutral pH, making it safe to use on most hard, non-porous surfaces. EPA Registration Number: 10324-154-34714.

WHY IS IT USED?

WINTAFECT has a broad spectrum of uses. Perfect for not only hospitals but also restaurants and bars, supermarkets, public facilities, hotels, kitchens, crime scenes and schools. WINTAFECT is a one-step product with no rinsing± required and deodorizing capabilities that destroy odor causing bacteria at the source.



HOW IS IT USED?

*It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **Read entire label before use.** Allow surfaces to come to room temperature (68 °F, 20 °C) before treatment with this product. Before using this product, food products and packaging materials must be removed from the room or carefully protected. This product is not for use on medical device surfaces.*

FOR USE AS A ONE-STEP DISINFECTANT/FUNGICIDE/VIRUCIDE*:

1. Pre-clean visibly soiled areas.
2. Apply use solution of 2 oz. of this product per gal. of water to disinfect hard, non-porous, non-food contact surfaces with a sponge, brush, cloth, mop, auto scrubber, mechanical spray device, hand pump trigger spray device, coarse trigger spray device. For spray applications, spray 6-8 inches from surface. Do not breathe spray.
3. Treated surfaces must remain visibly wet for 2 minutes to kill SARS-CoV-2 or for 10 minutes to kill all organisms listed on the label.
4. Allow to air dry. Rinsing of floors is not necessary unless they are to be waxed or polished.
5. Prepare a fresh solution daily or when visibly dirty.

BLOODBORNE PATHOGEN INSTRUCTIONS: *KILLS HIV, HBV AND HCV ON PRE-CLEANED HARD, NON-POROUS SURFACES/OBJECTS PREVIOUSLY OILED WITH BLOOD/BODY FLUIDS

in health care settings or other settings in which there is an expected likelihood of soiling of hard, non-porous surfaces/objects with blood or body fluids and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1 (HIV-1), Hepatitis B Virus (HBV) and Hepatitis C Virus (HCV).

±Food contact surfaces must be rinsed with potable water after application of product.

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1, HBV and HCV ON SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS:

- **Personal Protection:** Wear protective latex gloves, gowns, masks and eye protection.
- **Cleaning Procedure:** Blood and other body fluids containing HIV-1, HBV & HCV must be thoroughly cleaned from hard, non-porous surfaces/objects before application of this product.
- **Disposal of Infectious Materials:** Blood and other body fluids, cleaning materials and clothing must be autoclaved and disposed of according to federal, state and local regulations for infectious waste disposal.
- **Contact Time:** Allow hard, non-porous surfaces to remain visibly wet for 2 minutes to kill HIV and 10 minutes for all organisms listed on the label.

FOR USE AS A DISINFECTANT AND VIRUCIDE* BY ELECTROSTATIC SPRAYING: Before use, remove bystanders and pets from the area to be treated. Wear an N95 filtering facepiece or half face respirator, eye protection, and gloves when spraying.

1. Pre-clean visibly soiled surfaces.
2. Fill the sprayer with a use solution of 2 oz. of this product per gal. of water.
3. Set the unit to produce particles with a volume median diameter (VMD) of $\geq 40 \mu\text{m}$.
4. For units that allow the electrostatic spray function to be turned on or off, set the electrostatic spray function to ON.
5. Spray hard, non-porous surfaces from 6 - 18 inches away until thoroughly wet.
6. Allow surfaces to remain visibly wet for 10 minutes, reapplying if necessary. A 10-minute contact time is necessary to kill SARS-CoV-2 when Wintafect is applied by electrostatic sprayer.
7. Allow spray to settle before reentering treated areas.
8. Wipe dry with a clean cloth or allow to air dry.

TO CONTROL MOLD/MILDEW:

1. Pre-clean visibly soiled areas.
2. Prepare use solution by adding 2 oz. of this product per gal. of water.
3. Apply use solution to hard, non-porous surfaces which will effectively inhibit the growth of mold and mildew and their odors.
4. Repeat treatment every seven days, or more often if new growth appears.

TO CONTROL THE GROWTH OF MOLD AND MILDEW ON NON-POROUS ATHLETIC EQUIPMENT:

For use on wrestling and gymnastic mats, athletic mats, exercise equipment, athletic training tables, physical therapy tables, athletic helmets, wrestling/boxing headgear, athletic shoe soles, and other hard, non-porous surfaces. Thoroughly clean surfaces with soap or detergent and rinse with water. Prepare a use solution of 2 oz. of this product per gal. of water. Apply by sponge, brush, cloth, mop, mechanical spray device, hand pump trigger spray device. For spray applications, spray 6-8 inches from surface. Do not breathe spray. Ventilate buildings and other closed spaces. Do not use equipment until treatment has been absorbed, set or dried. Repeat treatment every seven days, or more often if new growth appears.



Solutions for a Cleaner Environment

#0900 | Wintafect Disinfectant, Cleaner, & Deodorizer

FOR USE AS A DISINFECTANT, FUNGICIDE, VIRUCIDE* OF GROOMING CLIPPERS, HAIR CLIPPERS, ELECTRIC SHEARS, BARBER/SALON SHEARS AND OTHER IMPLEMENTS, SALON INSTRUMENTS AND TOOLS: Remove hair, dandruff and dust particles prior to disinfecting the blades. While clipper/shear is running, hold in the downward position and spray with a use solution of 2 oz. of this product per gal. of water (or equivalent use dilution). Do not spray on the clipper case or drip into clipper housing. Immerse pre-cleaned barber/salon tools such as combs, brushes, razors, tweezers, manicure/pedicure tools, scissors, and other salon instruments and tools in a use solution of 2 oz. of this product per gal. of water for at least 10 minutes. Rinse thoroughly and dry before use. Prepare a fresh solution daily or when visibly dirty. **Note:** Plastic instruments can remain immersed until ready to use. Stainless steel shears and other metal instruments must be removed after 10 minutes, rinsed, dried and kept in a clean, non-contaminated receptacle. Prolonged soaking will cause damage to metal instruments.

NON-ACID TOILET BOWL AND URINAL DISINFECTANT DIRECTIONS:

Remove visible soil prior to disinfection.

From a Concentrate: Add 2 oz. of product directly into the water in the toilet bowl. Brush thoroughly over exposed surfaces and under the rim with a toilet brush, cloth, or sponge. Allow surfaces to remain visibly wet for 10 minutes and flush.

From a Use Solution: Empty water out of toilet bowl or urinal and apply use-solution prepared by adding 2 oz. of this product per gal. of water to exposed surfaces, including under the rim with a toilet brush, cloth, or sponge. Allow surfaces to remain visibly wet for 10 minutes and flush.

TO CLEAN WATERFREE/WATERLESS URINALS: Remove any debris from the urinal. Spray 0.5 to 1 oz. of use solution onto urinal surface. **To prepare use solution:** Add 2 oz. per gal. of water. DO NOT spray product directly onto cartridge. Wipe surface to clean. Change cartridge as needed. The unit is ready for use.

FOR USE AS A CLEANER ON FINISHED FLOORS: To limit gloss reduction, use 2- 4 oz. of this product per gal. of water. Apply with a damp mop or auto scrubber. Allow surface to air dry.

AUTOMATED DILUTION SYSTEM INSTRUCTIONS: Remove cap from empty container. Fill empty container with a freshly prepared use-solution. Replace cap. Place correct use-dilution label on newly filled container.

PACKAGING:

WINTAFECT is available in Closed Loop[‡], gallon, 4 – 1 gallon case, 5 gallon & 55 gallon containers.

[‡]ALCO's Closed Loop System is a non-spill, metered dispensing system easily fills up quarts or gallons and is safer for the user because it minimizes chemical handling. By dispensing the proper amount of product every time, the dilution ratio remains constant, eliminating product waste and the possibility of over or under diluting the chemical - saving time and money!

WHAT DOES IT KILL?

DISINFECTION PERFORMANCE: This product kills the following bacteria in 10 minutes at 2 oz. per gal. of 400 ppm hard water (660 ppm active) and 5% soil on hard, non-porous surfaces:

- | | | |
|--|--|--|
| ▪ <i>Acinetobacter baumannii</i> | ▪ <i>Klebsiella pneumoniae</i> | ▪ <i>Staphylococcus aureus</i> (CA-MRSA) |
| ▪ <i>Acinetobacter lwoffii</i> | (NDM-1) | ▪ <i>Staphylococcus aureus</i> (MRSA) |
| ▪ <i>Bordetella bronchiseptica</i> | ▪ <i>Listeria monocytogenes</i> | ▪ <i>Staphylococcus epidermidis</i> |
| ▪ <i>Chlamydia psittaci</i> | ▪ <i>Micrococcus luteus</i> | ▪ <i>Staphylococcus epidermidis</i> (Ampicillin, Cefazolin, Oxacillin, Penicillin Resistant) |
| ▪ <i>Citrobacter freundii</i> | ▪ <i>Mycoplasma gallisepticum</i> | ▪ <i>Streptococcus agalactiae</i> |
| ▪ <i>Enterobacter agglomerans</i> | ▪ <i>Pasteurella multocida</i> | ▪ <i>Staphylococcus haemolyticus</i> |
| ▪ <i>Klebsiella aerogenes</i> | ▪ <i>Proteus vulgaris</i> | ▪ <i>Streptococcus pneumoniae</i> (Penicillin Resistant) |
| ▪ <i>Enterobacter cloacae</i> | ▪ <i>Pseudomonas aeruginosa</i> | ▪ <i>Streptococcus pyogenes</i> |
| ▪ <i>Escherichia coli</i> | ▪ <i>Pseudomonas aeruginosa</i> (Tetracycline Resistant) | ▪ <i>Streptococcus mutans</i> |
| ▪ <i>Escherichia coli</i> O157:H7 | ▪ <i>Pseudomonas cepacia</i> | ▪ <i>Staphylococcus aureus</i> (VISA) |
| ▪ <i>Escherichia coli</i> (Carbapenem Resistant) | ▪ <i>Salmonella enterica</i> | ▪ <i>Vibrio cholerae</i> |
| ▪ <i>Escherichia coli</i> (ESBL) | ▪ <i>Salmonella enteritidis</i> | ▪ <i>Yersinia enterocolitica</i> |
| ▪ <i>Escherichia coli</i> (Tetracycline Resistant) | ▪ <i>Salmonella enterica</i> serotype pullorum | |
| ▪ <i>Enterococcus faecalis</i> (VRE) | ▪ <i>Salmonella typhi</i> | |
| ▪ <i>Enterococcus hirae</i> | ▪ <i>Salmonella typhimurium</i> | |
| ▪ <i>Fusobacterium necrophorum</i> | ▪ <i>Serratia marcescens</i> | |
| ▪ <i>Klebsiella oxytoca</i> | ▪ <i>Shigella flexneri</i> | |
| ▪ <i>Klebsiella pneumoniae</i> | ▪ <i>Shigella sonnei</i> | |
| | ▪ <i>Staphylococcus aureus</i> | |

VIRUCIDAL PERFORMANCE: This product kills the following viruses in 10 minutes at 2 oz. per gal. of 400 ppm hard water (660 ppm active) and 5% soil on hard, non-porous surfaces:

- | | | |
|--|--|--------------------------------------|
| ▪ <i>Avian Influenza A</i> (H5N1) | ▪ <i>Hepatitis C Virus</i> (Bovine Viral Diarrhea Virus) | ▪ <i>Influenza A</i> (H1N1) |
| ▪ <i>Avian Influenza A</i> (H3N2) | ▪ <i>Herpes Simplex Type 1</i> | ▪ <i>Influenza A</i> (H3N2) |
| ▪ <i>Cytomegalovirus</i> | ▪ <i>Herpes Simplex Type 2</i> | ▪ <i>Norovirus</i> |
| ▪ <i>Coronavirus</i> (SARS-associated) | ▪ <i>Human Coronavirus</i> | ▪ <i>Respiratory Syncytial Virus</i> |
| ▪ <i>Hepatitis B Virus</i> | ▪ <i>Human Immunodeficiency Virus Type 1</i> *** (HIV 1) | ▪ <i>SARS-CoV-2</i> *** |
| | | ▪ <i>Vaccinia Virus</i> |

*** Indicates a 2-minute contact time is required for this claim.

FUNGICIDAL PERFORMANCE: This product kills the following fungi in 10 minutes at 2 oz. per gal. of 400 ppm hard water (660 ppm active) and 5% soil on hard, non-porous surfaces:

- | | |
|---------------------------|--|
| ▪ <i>Candida albicans</i> | ▪ <i>Trichophyton interdigitale</i> (Athlete's foot) (Ringworm fungus) |
|---------------------------|--|

MILDEWSTATIC PERFORMANCE: This product controls the following mold at 2 oz. per gal. of 400 ppm hard water (660 ppm active) and 5% soil on hard, non-porous surfaces for up to 7 days:

- *Aspergillus niger*